



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let $\operatorname{sn} u = x$. $\therefore x^4 + 2x^2 + 1/\kappa^2 = 0$, where κ is the modulus.

$$\therefore (x^2 + 1)^2 = -(1 - \kappa^2)/\kappa^2.$$

$$\therefore x = \pm \sqrt{\frac{-\kappa \pm \sqrt{(-1)\sqrt{1-\kappa^2}}}{\kappa}} = \pm n, \text{ suppose. } \therefore \operatorname{sn} u = \pm n.$$

$$\begin{aligned} \therefore u &= 2 \int_0^n \frac{dx}{\sqrt{[(1-x^2)(1-\kappa^2 x^2)]}} = 2F(\kappa, n) \\ &= 2[n + \frac{1}{6}n^3(1+\kappa^2) + \frac{1}{40}n^5(3+2\kappa^2+3\kappa^4) + \dots]. \end{aligned}$$

When $\kappa=1$, $u=2\sqrt{-1}(1-\frac{1}{3}+\frac{1}{5}-\frac{1}{7}+\dots)$, since $n=\pm\sqrt{-1}$.

$$\therefore u = \frac{1}{2}\pi \sqrt{-1}.$$

$$\text{But } u = 2 \int_0^{\sqrt{-1}} \frac{dx}{1-x^2} = \log \left(\frac{1+\sqrt{-1}}{1-\sqrt{-1}} \right).$$

$$\therefore \pi = 2\sqrt{-1} \log \left(\frac{1-\sqrt{-1}}{1+\sqrt{-1}} \right), \text{ a result previously referred to in this journal.}$$

PROBLEMS FOR SOLUTION.

ARITHMETIC.

167. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

A traveler notices that $m=2\frac{1}{2}$ times the number of spaces between the telegraph poles that he passes in a minute is the rate of train in miles per hour. How far apart are the poles?

168. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

If I buy for $m=10\%$ and $n=5\%$ less I shall gain $p=15\%$ and $q=5\%$ more. What is my rate of gain?

ALGEBRA.

178. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

A_n being the arithmetic mean of the n th powers of the numbers less than p and prime to it, find a relation between A_3 , A_2 and p .

179. Proposed by DR. L. E. DICKSON, The University of Chicago.

Find the roots of the algebraically solvable quintic equation

$$x^5 + qx^2 + px + \frac{1}{5} \left(\frac{q^2}{p} - \frac{p^3}{5q} \right) = 0.$$